arXiv:1612.09567v2 [math.GN] 5 Mar 2017

The weak Hilbert-Smith conjecture from a Borsuk-Ulam-type conjecture

Alexandru Chirvasitu, Ludwik Dąbrowski, and Mariusz Tobolski

Abstract

We show that the Borsuk-Ulam-type conjecture of Baum, Dąbrowski and Hajac partially settles a conjecture of Ageev. In turn, the latter implies the weak version Hilbert-Smith conjecture stating that no infinite compact zero-dimensional group can act freely on a manifold such that the orbit space is finite-dimensional.

Key words: Hilbert-Smith conjecture, Ageev conjecture, Borsuk-Ulam theorem, p-adic integers, free action, Menger compactum, dimension

MSC 2010: 22C05; 54H15; 57S10

Introduction

The fifth question on the Hilbert's famous list of problems [10] concerned a characterization of Lie groups. In a modern and commonly accepted form [27, Theorem 2.7.1] the result reads as follows.

Theorem 1 *[Hilbert's fifth problem] Let G be a topological group which is locally Euclidean. Then G is isomorphic to a Lie group.*

The above theorem was proved by Gleason [9, Theorem 3.1] and by Montgomery and Zippin [20, Theorem A] in 1952. The theorem and the mathematical tools developed in the course of proving it have many applications, including the celebrated Gromov theorem on groups of polynomial growth (see [27] for a survey of applications).

It is worth noting that in Hilbert's lecture [10] groups were not treated abstractly, but rather as groups of transformations. Hence, one might argue that the following generalization of Theorem 1 in fact better captures the original intent behind Hilbert's problem (see e.g. [27, Conjecture 2.7.2]).

Conjecture 2 *[Hilbert-Smith conjecture] Let G be a locally compact topological group acting continuously and effectively on a connected finite-dimensional manifold X. Then, G is isomorphic to a Lie group.*

It is known that the Hilbert-Smith conjecture can be reduced to studying actions of the groups $\mathbb{Z}_p$ of p-adic integers for arbitrary primes p. This was proved by Lee [14, Theorem 3.1] using the Gleason-Yamabe theorem on the structure of locally compact groups [28, Theorem 5'], and some results of Newman [23]. We state the reduced conjecture following [27, Conjecture 2.7.3].

Conjecture 3 *[Hilbert-Smith conjecture for p-adic actions] It is not possible for a p-adic group $\mathbb{Z}_p$ to act continuously and effectively on a connected finite-dimensional manifold X.*

Let us recall some results that partially confirm the Hilbert-Smith conjecture. In [21] Montgomery and Zippin observed that Conjecture 2 is true for transitive actions on topological manifolds and for smooth actions on smooth manifolds. Repovš and Ščepin [26] proved it for Lipschitz actions. Then Martin announced the proof for quasiconformal actions on Riemannian manifolds in [16]. There is also a proof for Hölder actions given by Maleshich [15]. Recently, Pardon showed that Conjecture 2 is true for three-manifolds [24]. In the survey [7] Dranishnikov gives an account of various partial results and reduces a weaker version of the conjecture to two other problems. In its full generality, the Hilbert-Smith conjecture remains unsolved.

We base our approach to the Hilbert-Smith conjecture on classifying spaces for p-adic groups and the theory of Menger compacta [18, 4]. In [8] Dranishnikov shows that any zero-dimensional compact metric group (e.g. $\mathbb{Z}_p$) can act freely on universal Menger compacta. Such actions can be fairly exotic, e.g. for the n-dimensional Menger compactum μ^n the dimension of the orbit space $\mu^n/\mathbb{Z}_p$ can exceed n.

Nevertheless, there is a class of actions for which $\dim(\mu^n/\mathbb{Z}_p) = n$; these are all proven isomorphic (for given G, in the category of G-spaces) by Ageev in [2]. The results of the latter paper also suggest that the actions in question satisfy a universality condition to the effect that for any free action of G on an at-most-n-dimensional compact space Y, every equivariant map $Y \to \mu^n$ is approximable by an equivariant embedding. This is referred to as 'strong G-n universality' in the discussion preceding [2, Theorem B], and is an equivariant version of the celebrated Bestvina recognition criterion for Menger compacta ([4] and [2, Theorem A (b)]).

In [1] Ageev states a certain conjecture about Menger compacta which if true would imply a weaker version of the Hilbert-Smith conjecture for free actions on manifolds with finite-dimensional orbit spaces. The following statement appears as [1, Conjecture].

Conjecture 4 *[Ageev] Suppose two Menger compacta μ^m and μ^n with $m > n$ are acted on freely by a non-trivial zero-dimensional compact metric group G. Then, there is no equivariant map $f : \mu^m \to \mu^n$.*

These ideas were followed by Yang in [29], where a free Menger $\mathbb{Z}_p$-compactum is explicitly constructed.

Baum, Dąbrowski and Hajac [3, Conjecture 2.2] proposed the generalized compact-Hausdorff-space Borsuk-Ulam conjecture. In order to state it, recall first that for two spaces X and Y, their *topological join* $X * Y$ is the space defined by

$$X * Y = X \times Y \times [0,1]/\sim,$$

where the equivalence relation identifies all $(x, y_0, 0)$, $x \in X$ for fixed $y_0 \in Y$, and similarly identifies all $(x_0, y, 1)$, $y \in Y$ for fixed $x_0 \in X$.

Conjecture 5 *Let X be a compact Hausdorff space equipped with a continuous free action of a non-trivial compact Hausdorff group G. Then, for the diagonal action of G on the topological join $X * G$, there does not exist a G-equivariant continuous map $f : X * G \to X$.*

Here we show that the positive answer to Conjecture 5 would prove Conjecture 4 as well. This result would then be sufficient to resolve affirmatively the finite-dimensional-orbit-space version of the Hilbert-Smith conjecture for free actions in Corollary 15.

The proposed reformulation of the problem is justified by recent work on and interest in Borsuk-Ulam type theorems. Conjecture 5 is known to be true for actions of finite cyclic groups on compact Hausdorff spaces, which can be deduced from [17, Theorem 6.2.6] and was recently proved using

different methods by Passer in [25, Theorem 2.8]. Moreover, in [6, Section 3] Chirvasitu and Passer propose a possible approach to Conjecture 5 in its full generality. Let us notice that in [1] Ageev mentions that a Borsuk-Ulam type result of Edwards and Bestvina (unpublished), generalizing [17, Theorem 6.2.6] to zero-dimensional compact metric groups, speaks in favor of Conjecture 4.

For the convenience of the reader, we include below a diagrammatic depiction of the logical dependence of the results below. We denote conjectures, lemmas and theorems by C, T and L respectively, and corollaries by Cor.



1 Preliminaries

All of our topological spaces are at the very least second countable and metrizable, and *compacta* are compact Hausdorff metrizable topological spaces. We denote by μ^n the universal n-dimensional Menger compactum characterized abstractly in a number of ways in [4].

We use [13] as our source of background for the dimension theory of separable metrizable spaces. We will not recall any of the precise definitions of dimension here (henceforth denoted by dim), and refer instead e.g. to [13, Definition III 1]. Suffice it to say the concept captures the usual intuition, and specializes to the standard notion of dimension for manifolds. The universality of μ^n referred to above consists in the fact that it is n-dimensional, and contains a homeomorphic copy of every n-dimensional compactum.

Following [1, Definition 1], we recall

Definition 6 Let n be a non-negative integer and G a topological group acting freely on a space X. Then, X equipped with the action in question is *free n-universal* (or simply *universal* for short if n is understood) if for any free G-space Y with

$$\dim(Y/G) \leqslant n$$

and any closed G-invariant subspace $Z \subseteq Y$ every equivariant map $Z \to X$ admits an extension to an equivariant map $Y \to X$. ♦

In Section 2 below we will make crucial use of the following result (see [1, Theorem 1]).

Theorem 7 *A Menger compactum μ^n equipped with a free action by a zero-dimensional compact group is free n-universal in the sense of Definition 6.* ■

Remark 8 Note that although in the setting of the present paper our topological Hausdorff spaces are assumed to be metrizable, in Conjecture 5 the metrizability of spaces and groups is not required. ♦

We will need a few more scattered pieces of notation and terminology:

Notation 9 In the context of a group G acting on spaces X and Y, we write $X \times_\Delta Y$ (with the group and actions being understood) to denote the Cartesian product $X \times Y$ equipped with the diagonal G-action. ♦

Finally, we will on occasion refer to *ANR* spaces (for absolute neighborhood retract). These are spaces X with the property that whenever embedded homeomorphically $\iota : X \to Y$ as a closed subspace of a topological space Y, there is an open neighborhood U of X in Y that retracts onto X in the sense that we have a diagram of the form



for some $r : U \to X$ so that the X-based loop is the identity. Metrizable manifolds are automatically ANR, e.g. by [12, Theorem V.7.1] or the main result of [5].

2 Main results

We recall the following version of the Hilbert-Smith conjecture in its weak form, whereby free actions are substituted for effective ones, and an additional constraint is imposed on the orbit space. See e.g. [7, Conjecture 1.2]; we strengthen that statement somewhat by allowing our groups to be *locally* compact, so long as the action is well-behaved enough for the resulting orbit spaces to be locally compact Hausdorff spaces.

Conjecture 10 *Let G be a locally compact group acting freely and properly on a topological manifold M such that the orbit space M/G is finite dimensional. Then G is a Lie group.*

An examination of the argument presented in [27, §2.7.2] will show that the reduction of the general Hilbert-Smith conjecture to its p-adic version (Conjecture 3) also goes through in the present setting of free actions and finite-dimensional orbit spaces:

Conjecture 11 *Let p be a prime number and M a topological manifold. Then $\mathbb{Z}_p$ cannot act freely on M so that $M/\mathbb{Z}_p$ is finite-dimensional.*

We now make the following observation, announced above.

Lemma 12 *Conjecture 11 implies Conjecture 10.*

Proof The reduction of the full Hilbert-Smith conjecture to its p-adic version as described in [27, §2.7.2] proceeds by passing to a copy of $\mathbb{Z}_p \subseteq G$ embedded in the arbitrary locally compact group G. Freeness of the action is of course preserved by this passage, and only the implication

$$\dim(M/G) < \infty \Rightarrow \dim(M/\mathbb{Z}_p) < \infty \tag{2.1}$$

needs verification.

Since the action of G on M is free, G embeds into M as an orbit and is hence metrizable and finite-dimensional. The Gleason-Yamabe structure theorem on locally compact groups ([27, Theorem 1.1.17]) shows that G is an extension of a Lie group by a compact group K. Furthermore, it follows from the general structure theory of finite-dimensional compact groups ([11, Theorems 8.22 and 9.55]) that K is a successive extension of Lie and zero-dimensional compact groups. Finally, all extensions in sight are locally trivial fibrations by [22, Corollary 2].

It follows from the above remarks that the family of fibers of $M \to M/G$ is *equi-LC^d* for all $d \geqslant 0$ in the sense of [19, discussion preceding Theorem 1.2], and hence the assumed finite-dimensionality of M/G implies via [19, Theorem 1.2] that $M \to M/G$ is a locally trivial bundle.

The above remarks, together with the fact that $G \to G/\mathbb{Z}_p$ is a locally trivial bundle by [22, Corollary 2], imply that $M/\mathbb{Z}_p \to M/G$ is a locally trivial fibration and hence

$$\dim(M/\mathbb{Z}_p) \leqslant \dim(M/G) + \dim(G/\mathbb{Z}_p) = \dim(M/G) + \dim(G) < \infty.$$

This concludes the proof of the desired bound (2.1). ■

As per Lemma 12, our aim will be to show that Conjecture 5 implies Conjecture 11. We follow the plan laid out in [1], as we now recall. First, we paraphrase [1, Theorem 4] in a form that will suit us here. This slightly more precise form of the statement is easily extracted from the proof in loc. cit.

Theorem 13 *Let p be a prime number and let the p-adic group $\mathbb{Z}_p$ act freely on the Menger compacta μ^m and μ^n for some $m > n$ so that*

$$\dim(\mu^m/\mathbb{Z}_p) < \infty.$$

Suppose furthermore that there are no $\mathbb{Z}_p$-equivariant maps $\mu^m \to \mu^n$.

Then, for any ANR space M acted upon freely by $\mathbb{Z}_p$ we have $\dim(M/\mathbb{Z}_p) > n$. ■

Given Theorem 13, [1] proposes to attack Conjecture 11 via Conjecture 4. However, somewhat weaker version of the latter is sufficient for our purposes:

Theorem 14 *Let $m > n$ be non-negative integers, and let the zero-dimensional non-trivial compact group G act freely on μ^m and μ^n so that $\dim(\mu^n/G) \leqslant m$. Suppose that there are no G-equivariant maps $\mu^n * G \to \mu^n$.*

Then, there are no G-equivariant maps $\mu^m \to \mu^n$.

Notice that Theorem 14 assumes that Conjecture 5 is satisfied for a zero-dimensional compact metric group acting on a compact Hausdorff space of finite covering dimension. Before going into the proof, let us record the resolution of the weak Hilbert-Smith conjecture.

Corollary 15 *Conjecture 5 implies Conjecture 11.*

Proof Let $m > n$ be arbitrary positive integers.

We first apply Theorem 14 to actions of $\mathbb{Z}_p$ on μ^m and μ^n constructed as in [8], having the properties

$$\dim(\mu^m/\mathbb{Z}_p) = m, \ \dim(\mu^n/\mathbb{Z}_p) = n.$$

The non-existence of an equivariant map $\mu^m \to \mu^n$ resulting from this theorem then ensures that the hypotheses of Theorem 13 are satisfied for this specific choice of free actions, and hence $\dim(M/\mathbb{Z}_p) > n$. Since n was arbitrary, this concludes the proof that the dimension of the orbit space cannot be finite. ■

Proof of Theorem 14 The freeness of all actions in sight imply that all orbits are equivariantly homeomorphic to G itself.

Now, our assumption that $\dim(\mu^n/G) \leqslant m$ shows via Corollary 17 below that we have

$$\dim(\mu^n * G/G) \leqslant m. \tag{2.2}$$

The universality of the G-action on μ^m provided by Theorem 7 and (2.2) ensure that any equivariant homeomorphism from a G-orbit in $\mu^n * G$ onto a G-orbit in μ^m extends to an equivariant map

$$\mu^n * G \to \mu^m.$$

Composition with an equivariant map $\mu^m \to \mu^n$ as in the statement of the theorem would then imply the existence of an equivariant map $\mu^n * G \to \mu^n$, contradicting the assumption. ■

Lemma 16 *Let X be a compact space equipped with a free action by a compact group G. Then, we have the inequality*

$$\dim(X * G/G) \leqslant \max(\dim(X/G), \dim(X) + 1). \tag{2.3}$$

Proof The orbit space $X*G/G$ on the left hand side of (2.3) is the union of the two closed subspaces X/G (at the endpoint $1 \in [0,1]$; see the definition of joins in the introduction) and $\{*\} \cong G/G$ at $0 \in [0,1]$, and the open subspace

$$X \times_\Delta G \times J/G \cong (X \times_\Delta G/G) \times J$$

where $J = (0,1)$ and the Δ subscript is explained in Notation 9.

[13, Corollary 1 to Theorem III 2] implies the estimate

$$\dim(X * G/G) \leqslant \max(\dim(X/G), \dim((X \times_\Delta G/G) \times J)).$$

Hence, for our purposes, it suffices to prove that we have

$$\dim((X \times_\Delta G/G) \times J) \leqslant \dim(X) + 1. \tag{2.4}$$

First, notice that the subadditivity of dimension under taking Cartesian products ([13, Theorem III 4]) together with $\dim(J) = 1$ bounds the left hand side of (2.4) above by

$$\dim(X \times_\Delta G/G) + 1.$$

Finally, the desired conclusion follows from the fact that

$$(x, g) \mapsto (xg, g)$$

implements an isomorphism of G-spaces between $X \times G$ with the right hand factor action and $X \times_\Delta G$. This, in turn, ensures

$$X \times_\Delta G/G \cong X,$$

and hence

$$\dim(X \times_\Delta G/G) = \dim(X).$$

This finishes the proof of the lemma. ■

As an immediate consequence of Lemma 16, we obtain

Corollary 17 *Let $m > n$ be non-negative integers. For a free action of a compact group G on μ^n with $\dim(\mu^n/G) \leqslant m$ we have $\dim(\mu^n * G/G) \leqslant m$.* ■

Remark 18 Theorems 13 and 14 are stated in such generality to emphasize their connection with Ageev's original formulation [1]. However, working with the universal actions on Menger compacta μ^n such that $\dim(\mu^n/\mathbb{Z}_p) = n$ would still confirm Conjecture 11.

As mentioned in the introduction, the actions in question are proved unique up to isomorphism (for given n) in [2, Theorem B] and their existence is proven by [8, Theorem 1]. Remark 1 on p. 228 therein argues that the orbit space for the action on μ^n being constructed in the proof of the theorem is indeed n. ♦

Acknowledgements

We acknowledge a suggestion by A. Volovikov that the conjecture made in [3] should prove Conjecture 4, and discussions with H. Toruńczyk and with S. Spież, who suggested using the universality of Menger compacta. We thank R. Edwards for confirmation of his unpublished results mentioned in the paper of S. M. Ageev. We are also grateful to Piotr M. Hajac and Benjamin Passer for numerous conversations on the contents of this paper and related topics contributing considerably to its improvement.

This work is part of the project Quantum Dynamics sponsored by EU-grant RISE 691246 and Polish Government grant 317281. A.C. was partially supported by NSF grant DMS-1565226. L.D. is grateful for his support at IMPAN provided by Simons-Foundation grant 346300 and a Polish Government MNiSW 2015–2019 matching fund. Finally, M.T. was partially supported by the project Diamentowy Grant No. DI2015 006945 financed by the Polish Ministry of Science and Higher Education.

References

[1] S. M. Ageev. Classifying spaces for free actions and the Hilbert-Smith conjecture. *Mat. Sb.*, 183(1):143–151, 1992.

[2] S. M. Ageev. Characterization of the free action of a zero-dimensional compact group on a k-dimensional Menger compactum. *Izv. Ross. Akad. Nauk Ser. Mat.*, 59(2):3–46, 1995.

[3] Paul F. Baum, Ludwik Dąbrowski, and Piotr M. Hajac. Noncommutative Borsuk-Ulam-type conjectures. *Banach Center Publ.*, 106:9–18, 2015.

[4] Mladen Bestvina. Characterizing k-dimensional universal Menger compacta. *Mem. Amer. Math. Soc.*, 71(380):vi+110, 1988.

[5] Robert Cauty. Une caractérisation des rétractes absolus de voisinage. *Fund. Math.*, 144(1):11–22, 1994.

[6] A. Chirvasitu and B. Passer. Compact Group Actions on Topological and Noncommutative Joins. *ArXiv e-prints*, April 2016.

[7] A. Dranishnikov. On some problems related to the Hilbert-Smith conjecture. *ArXiv e-prints*, April 2015.

[8] A. N. Dranishnikov. Free actions of 0-dimensional compact groups. *Izv. Akad. Nauk SSSR Ser. Mat.*, 52(1):212–228, 1988.

[9] Andrew M. Gleason. Groups without small subgroups. *Ann. of Math. (2)*, 56:193–212, 1952.

[10] David Hilbert. Mathematical problems. *Bull. Amer. Math. Soc. (N.S.)*, 37(4):407–436 (electronic), 2000. Reprinted from Bull. Amer. Math. Soc. **8** (1902), 437–479.

[11] Karl H. Hofmann and Sidney A. Morris. *The structure of compact groups*, volume 25 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, 2013. A primer for the student—a handbook for the expert, Third edition, revised and augmented.

[12] Sze-tsen Hu. *Theory of retracts*. Wayne State University Press, Detroit, 1965.

[13] Witold Hurewicz and Henry Wallman. *Dimension Theory*. Princeton Mathematical Series, v. 4. Princeton University Press, Princeton, N. J., 1941.

[14] Joo Sung Lee. Totally disconnected groups, p-adic groups and the Hilbert-Smith conjecture. *Commun. Korean Math. Soc.*, 12(3):691–699, 1997.

[15] Ĭozhe Maleshich. The Hilbert-Smith conjecture for Hölder actions. *Uspekhi Mat. Nauk*, 52(2(314)):173–174, 1997.

[16] Gaven J. Martin. The Hilbert-Smith conjecture for quasiconformal actions. *Electron. Res. Announc. Amer. Math. Soc.*, 5:66–70 (electronic), 1999.

[17] Jiří Matoušek. *Using the Borsuk-Ulam theorem*. Universitext. Springer-Verlag, Berlin, 2003. Lectures on topological methods in combinatorics and geometry, Written in cooperation with Anders Björner and Günter M. Ziegler.

[18] Karl Menger. *Kurventheorie*. Herausgegeben unter Mitarbeit von Georg Nöbeling. Second edition. Chelsea Publishing Co., Bronx, N.Y., 1967.

[19] Ernest Michael. Continuous selections. II. *Ann. of Math. (2)*, 64:562–580, 1956.

[20] Deane Montgomery and Leo Zippin. Small subgroups of finite-dimensional groups. *Ann. of Math. (2)*, 56:213–241, 1952.

[21] Deane Montgomery and Leo Zippin. *Topological transformation groups*. Robert E. Krieger Publishing Co., Huntington, N.Y., 1974. Reprint of the 1955 original.

[22] Paul S. Mostert. Local cross sections in locally compact groups. *Proc. Amer. Math. Soc.*, 4:645–649, 1953.

[23] M. H. A. Newman. A theorem on periodic transformations of spaces. *The Quarterly Journal of Mathematics*, (1):1–8, 1931.

[24] John Pardon. The Hilbert-Smith conjecture for three-manifolds. *J. Amer. Math. Soc.*, 26(3):879–899, 2013.

[25] Benjamin Passer. Saturated Actions on C*-algebra Suspensions and Joins by Finite Cyclic Groups. *To appear in IUMJ.*

[26] Dušan Repovš and Evgenij Ščepin. A proof of the Hilbert-Smith conjecture for actions by Lipschitz maps. *Math. Ann.*, 308(2):361–364, 1997.

[27] Terence Tao. *Hilbert's fifth problem and related topics*, volume 153 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2014.

[28] Hidehiko Yamabe. A generalization of a theorem of Gleason. *Ann. of Math. (2)*, 58:351–365, 1953.

[29] Zhiqing Yang. A construction of classifying spaces for p-adic group actions. *Topology Appl.*, 153(1):161–170, 2005.

Department of Mathematics, University of Washington, Seattle, WA 98195-4350, USA
E-mail address: chirva@uw.edu

SISSA (Scuola Internazionale Superiore di Studi Avanzati), Via Bonomea 265, 34136 Trieste, Italy
E-mail address: dabrow@sissa.it

Instytut Matematyczny, Polska Akademia Nauk, ul. Śniadeckich 8, 00-656 Warszawa, Poland
E-mail address: mtobolski@impan.pl